

Assay specificity:

Monoclonal Antibody

Mold Species


	<i>Aspergillus versicolor</i>	<i>Stachybotrys chartarum</i>	<i>Aspergillus fumigatus</i>	<i>Penicillium chrysogenum</i>	<i>Penicillium expansum</i>	<i>Trichoderma harzianum</i>	<i>Chaetomium globosum</i>
5F5	1.38	<0.11	<0.11	<0.11	<0.11	<0.11	<0.11
9A5	2.26	<0.11	<0.11	<0.11	<0.11	<0.11	<0.11

Values are O.D. in ELISA (405 nm, assay background <0.11)



AveX ELISA kit (5F5/9A5)
For *Aspergillus versicolor*
Antigen Detection
Product Code: EL-AVX
Lot: XXXXX

Sample Curve:



Content:

- Vial 1 (red top) 100 µl
Monoclonal antibody 5F5
- Vial 2 (white top) 400 µl
Aspergillus versicolor Standard
Concentration: 10,000U/ml
- Vial 3 (brown) 100 µl
Biotinylated monoclonal antibody 9A5
Dilute: 1:1000 for use

Storage: The ELISA kit should be stored at 4°C

For research and commercial use in vitro: not for human in vivo or therapeutic use.

An InBio™ product



www.inbio.com

Indoor Biotechnologies, Inc.
700 Harris Street,
Charlottesville, VA 22903
United States

Indoor Biotechnologies Ltd.
Vision Court
Caxton Place
Cardiff, Wales CF23 8HA
United Kingdom

Indoor Biotechnologies,
India Private Limited
Bangalore Bioinnovation Centre,
BioTech Park, Electronic City Phase 1
Bangalore-560100, India

Tel: (434) 984-2304
Fax: (434) 984-2709
Email: mail@inbio.com

Tel: +44 (0) 29 2167 4640
Email: Info@indoorbiotech.co.uk

Tel: +91-9901722009
Email: info@inbioindia.com

Certificate of Analysis

Monoclonal Antibody: 5F5 (clone 5F5 H2 H8)
Product Code: MA-5F5
Immunogen: *Aspergillus versicolor* extract
Isotype: Mouse IgG2a
Specificity: Binds to a species specific epitope of *Aspergillus versicolor* antigen.
Purification: Produced in tissue culture and purified by chromatography using Protein A. Single heavy and light chain bands on SDS-PAGE.
Concentration: 1 mg/ml in phosphate buffered saline, pH 7.4. Based on A280 for IgG (1.42=1mg/ml) 0.22µm filtered, preservative free.
Lot Number: xxxxx

Monoclonal Antibody: 9A5 (clone 9A5 B9 C10)
Product Code: BI-9A5
Immunogen: *Aspergillus versicolor* extract
Isotype: Mouse IgG2a
Specificity: Binds to a species specific epitope of *Aspergillus versicolor* antigen.
Purification: Produced in tissue culture and purified by chromatography using Protein A. Single heavy and light chain bands on SDS-PAGE.
Biotinylation: Biotinylated and titrated for use in ELISA at 1/1000 dilution. Prepared in 1% BSA/50% glycerol/PBS, pH 7.4, 0.22µm filtered, preservative free.
Lot Number: xxxxx

Antigen Standard: *Aspergillus versicolor*
Product Code: ST-AV1
Composition: *Aspergillus versicolor* source material extract prepared in 1% BSA/50% glycerol/PBS, 0.22µm filtered, preservative free, pH 7.4
Concentration: 10,000U/ml AveX
Calibration: The concentration of the extract was measured using a total protein assay. The extract is calibrated in arbitrary units where 1 unit of AveX equals approximately 1 nanogram of total protein.
Lot Number xxxxx

ELISA protocol for AveX antigen.

1. Coat polystyrene microtiter plates (NUNC Maxisorp Cert. NUNC catalog # 439454) with 100µl mAb 5F5 at 10µl/10ml, i.e. 1/1000 dilution of stock, in 50mM carbonate-bicarbonate buffer, pH 9.6, incubate overnight at 4°C.
2. Wash wells 3x with PBS-0.05% Tween 20, pH 7.4 (PBS-T). Incubate for 30 min. at room temperature with 100µl/well of 1% BSA, PBS-T. Wash 3x with PBS-T.
3. Use doubling dilutions of the antigen standard to make a control curve ranging from 1000 - 2U/ml AveX: Pipette 20µl of the standard into 180µl 1% BSA, PBS-T into wells A1 and B1 on the ELISA plate. Mix well and transfer 100µl across the plate into 100µl 1% BSA, PBS-T diluent to make 10 serial doubling dilutions. Wells A11, B11 and A12, B12 should contain only 1% BSA, PBS-T as blanks.
4. Add 100µl of diluted allergen samples and incubate for 1 hour at room temperature. Environmental samples for AveX analysis are routinely diluted two-fold from 1/2-1/8 for dust and air filter extracts or 1/10-1/80 for swabs, wallboard, and culture extracts.
5. Wash wells 3x with PBS-T and add 100µl diluted biotinylated anti-AveX mAb 9A5. The antibody solution contains 50% glycerol and should be diluted 1/1000 (i.e. 10µl/10ml) in 1% BSA, PBS-T. Incubate for 1 hour at room temperature.
6. Wash wells 3x with PBS-T and add 100µl diluted Streptavidin-Peroxidase (Sigma S5512, 0.25mg reconstituted in 1 ml distilled water). The reconstituted Streptavidin should be diluted 1/1000 (i.e. 10µl/10ml) in 1%BSA, PBS-T. Incubate for 30 minutes at room temperature.
7. Wash wells 3x with PBS-T and develop the assays by adding 100µl 1mM ABTS in 70mM citrate phosphate buffer, pH 4.2 containing a 1/1000 dilution of 30% H₂O₂ (i.e. 10µl/10ml ABTS). Read the plate when the optical density at 405nm reaches 2.0-2.4.

Notes:

The antigen standard is recommended for immunoassay calibration purposes only. Not recommended for in-vitro antibody measurements, T cell studies, immunization purposes, or other uses.

Buffer recipes, storage conditions and a list of frequently asked questions can be found under "Protocols" on our web site: www.inbio.com.

For research and commercial use in vitro: not for human in vivo or therapeutic use.